Culture Information

Sweet Pea

Sweet Pea seed can be poisonous if swallowed.

Plant type: Annual

Sowing: Sow Autumn and Spring.

Use a low nutrient sowing compost. Sweet Pea seedlings are sensitive

to high nutrient concentration. Temperature: 20°C at seed level.

Cover with thick layer (7 - 10 mm) of medium grade Vermiculite. Germinate in light (we use normal light conditions of 16 hours light).

Seedling emergence 1 - 2 weeks.

Various techniques of pre-soaking seed or chipping the seed coat have been suggested to improve germination. We have no evidence to

indicate that either of these techniques is of use.

Damping off can occur with Sweet Pea seedlings. Do not over water

emerging seedlings.

Growing on:

Temperature should be lowered to no more than 10°C once the

seedlings have emerged.

pH 6.5

Feeding Plants should not be overfed with nitrogen. Many growers find a high

potassium compound fertiliser suitable for the crop.

Growth regulation Use of growth regulants can increase branching (though not essential).

Pack/Pot size Traditionally 10 – 12 seeds in a 9cm pot.

Advice It is essential to maintain good moisture throughout production.

Plants should never dry out, but some air is needed in the substrate as well – it should not be heavy and saturated. A balance is needed.

Common diseases Botrytis, Mildews

Ball Colegrave cultural information is issued as a guide to growers, based on our own trials experience. It is not intended as a blueprint for growing. Any chemicals referred to should be used only in accordance with the manufacturer's instructions.